



04/10/98

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

ATTORNEYS AT LAW

SUITE 600

1100 NEW YORK AVENUE, N.W.

WASHINGTON, D.C. 20005-3934

(202) 371-2600

FACSIMILE (202) 371-2540 (202) 371-6566

ROBERT GREENE STERNE
EDWARD J. KESSLER
JORGE A. GOLDSTEIN
SAMUEL L. FOX
DAVID K.S. CORNWELL
ROBERT W. ESMOND
TRACY-GENE G. DURKIN
MICHELE A. CIMBALA
MICHAEL B. RAY
ROBERT E. SOKOHL
ERIC K. STEFFE
MICHAEL O. LEE

JOHN M. COVERT*
LINDA E. ALCORN
RAZ E. FLESHNER
ROBERT C. MILLONIG
STEVEN R. LUDWIG
MICHAEL V. MESSINGER
JUDITH U. KIM*
KEITH KIND
TIMOTHY J. SHEA, JR.
DONALD R. MCPHAIL
PATRICK E. GARRETT
BARBARA A. PARVIS
MICHAEL A. RAHMAN*

STEPHEN G. WHITESIDE*
NOEL B. WHITLEY*
JEFFREY T. HELVEY*
RICHARD A. DUNNING, JR.
KIMBERLIN M. TOOHEY
RALPH P. ALBRECHT
HEIDI L. KRAUS*
JEFFREY R. KURIN*
CARL B. MASSEY, JR.*
RAYMOND MILLIEN*
PATRICK D. O'BRIEN*
BRIAN S. ROSENBLUM*

DONALD J. FEATHERSTONE**
LAWRENCE B. BUGAISKY**
KAREN R. MARKOWICZ**
GRANT E. REED**
VICTOR E. JOHNSON**
SERGE SIRA**

*BAR OTHER THAN D.C.
**REGISTERED PATENT AGENTS

WRITER'S DIRECT NUMBER

April 10, 1998

INTERNET ADDRESS

Box Patent Application

Assistant Commissioner for Patents
Washington, D.C. 20231

Re: U.S. Non-Provisional Utility Patent Application under 37 C.F.R. § 1.53(b)
Appl. No. (to be assigned); filed (herewith)
For: **RNA Isolation Reagent and Methods**
Inventor: Domenica A. Simms
Our Ref: 0942.3840001/RWE

Sir:

The following documents are forwarded herewith for appropriate action by the U.S.
Patent and Trademark Office:

1. Utility Patent Application Transmittal (Form PTO/SB/05);
2. U.S. Utility Patent Application entitled:

RNA Isolation Reagent and Methods

and naming as inventor:

Domenica A. Simms

Assistant Commissioner for Patents
April 10, 1998
Page 2

the application comprising:


- 13 pages of description prior to the claims;
 - 4 pages of claims (claims 1-20); and
 - 1 page abstract;
3. Authorization to Treat a Reply As Incorporating An Extension of Time Under 37 C.F.R. § 1.136(a)(3) (in duplicate); and
4. Two (2) return postcards.

It is respectfully requested that, of the two attached postcards, one be stamped with the filing date of these documents and returned to our courier, and the other, prepaid postcard, be stamped with the filing date and unofficial application number and returned as soon as possible.

This patent application is being submitted under 37 C.F.R. § 1.53(b) without Declaration and without filing fee.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.



Robert W. Esmond
Attorney for Applicant
Registration No. 32,893

RWE/dab
P:\USERS\DENISEB\0942\3841cov
Enclosures

Please type a sign(+) inside this box



PTO/SB/05 (2/98)

Approved for use through 09/30/2000 OMB 0651-0032

Patent and Trademark Office U S DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

UTILITY PATENT APPLICATION TRANSMITTAL (Only for new nonprovisional applications under 37 CFR § 1.53(b))	Attorney Docket No	0942.3840001/RWE
	First Named Inventor or Application Identifier	Domenica A Simms
	Title	RNA Isolation Reagent and Methods
	Express Mail Label No	

APPLICATION ELEMENTS See MPEP chapter 600 concerning utility patent application contents	ADDRESS TO Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
--	--

1. ☐ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 18]
(preferred arrangement set forth below)
- Descriptive title of the invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R & D
- Reference to Microfiche Appendix
- Background of the invention
- Brief Summary of the invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure
6. ☐ Microfiche Computer Program (Appendix)
7. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
a. ☐ Computer Readable Copy
b. ☐ Paper Copy (identical to computer copy)
c. ☐ Statement verifying identity of above copies

3. ☐ Drawing(s) (35 U.S.C. 113) [Total Sheets]
4. ☐ Oath or Declaration [Total Pages]
a. ☐ Newly executed (original or copy)
b. ☐ Copy from a prior application (37 CFR 1.63(d)) (for continuation divisional with Box 17 completed) [Note Box 5 below]
i. ☐ **DELETION OF INVENTOR(S)**
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR §§ 1.63(d)(2) and 1.33(b)
5. ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein

ACCOMPANYING APPLICATION PARTS	
8. <input type="checkbox"/> Assignment Papers (cover sheet & document(s))	
9. <input type="checkbox"/> 37 CFR 3.73(b) Statement (when there is an assignee)	<input type="checkbox"/> Power of Attorney
10. <input type="checkbox"/> English Translation Document (if applicable)	
11. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449	<input type="checkbox"/> Copies of IDS Citations
12. <input type="checkbox"/> Preliminary Amendment	
13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) (Should be specifically itemized)	
14. <input type="checkbox"/> *Small Entity Statement(s)	<input type="checkbox"/> Statement filed in prior application. Status still proper and desired
15. <input type="checkbox"/> Certified Copy of Priority Document(s) (if foreign priority is claimed)	
16. <input checked="" type="checkbox"/> Other 37 CFR § 1.136(a)(3) Authorization	
<input type="checkbox"/> Other	
*NOTE FOR ITEMS 1 & 14: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 CFR § 1.27). EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 CFR § 1.28)	

17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment
- ☐ Continuation ☐ Divisional ☐ Continuation-in-Part (CIP) of prior application No: ____/____
- Prior application information Examiner _____ Group/Art Unit: _____

18. CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number or Bar Code Label [] or <input checked="" type="checkbox"/> Correspondence address below					
(Insert Customer No. or Attach bar code label here)					
NAME	STERNE, KESSLER, GOLDSTEIN & FOX P L L C				
	Attorneys at Law				
ADDRESS	Suite 600, 1100 New York Avenue, N W				
CITY	Washington	STATE	DC	ZIP CODE	20005-3934
COUNTRY	USA	TELEPHONE	(202) 371-2600	FAX	(202) 371-2540

NAME (Print/Type)	Robert W Esmond	Registration No. (Attorney Agent)	32,893
SIGNATURE		Date	April 10, 1998

Burden Hour Statement: this form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231

RNA Isolation Reagent and Methods

Background of the Invention

Field of the Invention

5 The present invention is in the field of molecular biology. The invention relates to improved methods, extraction reagents and kits for isolating RNA from eukaryotic cells, such as plant or animal cells.

Related Art

10 RNA isolation reagents and methods have been developed for isolating RNA, preferably cytoplasmic RNA and mRNA, from animal cells. These reagents and methods, when applied to the isolation of RNA from plant materials and plant cells, suffer from problems involving: the presence of Mg^{++} ions which degrade RNA; co-isolation of polysaccharides with RNA when strong chaotropic salts or ionic detergents are used; and the use of high salt concentrations requiring additional precipitation/redissolving steps. These methods give unsatisfactory results when applied to the isolation of RNA from plants.

15 U.S. Patent Numbers 5,346,994 and 4,843,155 disclose the use of extraction reagents having chaotropic salts (e.g., guanidinium isothiocyanate and ammonium isothiocyanate in high concentration (0.5-3M)), which completely disrupts cells and their nuclei, releasing RNA, DNA, proteins, membranous materials, and soluble polysaccharides into the extraction media. By virtue of the acidity and high salt concentration of the extraction media, the DNA precipitates out and is removed along with any insoluble cell debris during the centrifugation step.

20 Similar protocols are disclosed in Lewin, B.M., *Genes III*, John Wiley & Sons, publishers, New York, N.Y. (1990); Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second edition, Cold Spring Harbor

25

Laboratory, Cold Spring Harbor, NY (1989); and Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience, N.Y. (1987-1996).

5 The presence of chlorophyll and increased amounts of polysaccharides in plants provides additional problems for RNA isolation from such samples. In particular, the conventional use of chaotropic agents results in co-isolation of polysaccharides with the purified RNA. As plant cells have higher concentrations of polysaccharides, this problem is exacerbated by using known RNA isolation methods.

10 Furthermore, chlorophyll is broken down when plant cells are subjected to known RNA isolation methods and Mg^{++} ions are released from the chlorophyll. The Mg^{++} ions react with RNA and degrade it, reducing the yield and at least partially destroying the integrity of the RNA sequence.

15 Sambrook (*supra*, at sec. 7.6-7.9) discloses an alternative protocol for the isolation of cytoplasmic RNA from mammalian cells, which uses a non-disruptive RNA extraction reagent at physiological pH and salt concentration, instead of high concentrations of chaotropic salts. The extraction reagent contains an RNase inhibitor to protect the RNA during the isolation procedure and a nonionic detergent that solubilizes the cell membrane while leaving the nuclear membrane intact, to release cytoplasmic RNA. By not
20 disrupting the nucleus and other cell organelles, the cytoplasmic RNA in the extraction media is selectively extracted from the cell, and separated from cell debris by a centrifugation step. However, the Sambrook protocol suffers from the problem of requiring expensive RNase inhibitors and extensive sample handling to recover the purified mRNA. The presence of a cell wall was also a deterrent
25 from applying this technology to plant specimens.

Accordingly, improved methods and reagents are needed for RNA isolation from plants and plant or animal cells, which methods and reagents overcome one or more problems of known methods and reagents.

Summary of the Invention

The invention provides methods, extraction reagents and kits for RNA isolation from eukaryotic cells provided in a sample, where the use of at least a phenol, a chelator and a nonionic detergent replaces chaotropes and/or RNase inhibitors in the extraction reagent. These methods, reagents and kits provide superior results for isolation of cytoplasmic RNA or mRNA from eukaryotic cells, especially from plants, maintaining the integrity of the RNA without co-isolation of polysaccharides.

In a method of the present invention, an extraction reagent is used on fresh or frozen eukaryotic cells, or preferably frozen, powdered plant and animal samples. The RNA is localized in the aqueous phase after phase separation, and then precipitated with alcohol.

The process can comprise:

- (a) mixing the sample (or cell pellet) with the extraction reagent to form a mixture;
- (b) adding a haloalkane to the mixture and mixing;
- (c) separating the organic and aqueous phases by centrifugation or other known methods and transferring the aqueous phase; and
- (d) precipitating the cytoplasmic RNA from the aqueous phase obtained in step (c) with alcohol.

The process can optionally further comprise: (e) recovering the cytoplasmic RNA from the precipitate obtained in step (d). The process can also optionally further comprise: (f) isolating mRNA from the recovered cytoplasmic RNA (*e.g.*, as a precipitate) using any known method, such as by chromatography on oligo(dT)-cellulose (*see, e.g.*, Sambrook, *infra*, at §§ 7.26-7.29).

The extraction reagent preferably comprises:

- (a) at least one nonionic detergent (0.1-1.0% by volume);
- (b) at least one chelator (0.02-0.25 M);
- (c) at least one phenol (10%-60% by weight); and
- (d) at least one phenol solubilizer (15%-55% by volume).

The extraction reagent optionally further comprises:

- (e) at least one phenol stabilizer (0.05%-0.2% by weight);

The methods are useful for providing cytoplasmic or mRNA from cells contained in samples from eukaryotic organisms, with improved quantitative and/or qualitative yield over known methods, with RNA isolation from plants or plant cells preferred.

Another embodiment of the present invention relates to a kit comprising a carrier or receptacle being compartmentalized to receive and hold therein at least one container, wherein a first container contains at least one RNA reagent of the present invention, as described herein.

Other objects of the invention will be apparent to one of ordinary skill in the art from the following detailed description and examples relating to the present invention.

Detailed Description of the Preferred Embodiments

The invention is directed to methods and reagents using at least one of a phenol, a phenol stabilizer, a phenol solubilizer, a chelator and a nonionic detergent in the extraction reagent, in order to replace chaotropes and/or RNase inhibitors, and to provide enhanced isolation of intact cytoplasmic RNA from samples of eukaryotic organisms, tissues or cells.

In the present invention, subsequent sample handling is reduced substantially, as well as the material and labor costs for performing the RNA isolation. The use of a phenol, a chelator, a nonionic detergent, low salt concentrations, and no chaotropic agents improves the RNA isolation procedure, quantitatively and/or qualitatively.

The present invention provides several improvements over the related art, such as, but not limited to, at least one of: (a) the novel use of a chelator to protect the extracted RNA from degradation by Mg^{++} ions (*e.g.*, as present in animal cells, and as releasable from plant chlorophyll during the isolation process) in combination with phenol to protect the RNA from RNases; (b) the absence of

strong chaotropic agents from the RNA extraction media minimizes co-isolation of polysaccharides with the purified RNA; (c) the use of the low salt concentration in the RNA extraction medium, where (after addition of chloroform and phase separation) mRNA can be selected directly from the aqueous phase, while reducing or eliminating additional precipitation/redissolving steps.

The methods and reagents of the invention provide isolation of cytoplasmic RNA from eukaryotic organisms, such as, but not limited to, one or more of plants or plant materials or cells, animal cells or tissue, insects or insect cells, fungal hyphae or cells and other nucleated cells.

In a preferred embodiment, the eukaryotic sample (e.g., a plant sample) is preferably ground to a powder in liquid nitrogen, or by any other known method, where the sample remains frozen throughout the grinding procedure before RNA isolation. Alternatively, cell samples can be fresh or frozen but need not be ground into a powder. The invention thus provides direct isolation of cytoplasmic RNA or mRNA from eukaryotic cells.

RNA Extraction Reagents

In a preferred embodiment, the extraction reagent comprises:

- (a) at least one nonionic detergent (0.1-1.0% by volume) (e.g., tert-octylphenoxypoly(oxyethylene)ethanol (IGEPAL CA-630, Rhône Poulenc, France, 0.3-0.7%);
- (b) at least one chelator (0.02-0.25 M) (e.g., 0.05-0.5 M sodium citrate); and
- (c) at least one phenol (10%-60% by weight) (e.g., 20-40%); and
- (d) at least one phenol solubilizer (15%-55% by volume) (e.g., ethylene glycol, 22%).

The composition optionally further comprises:

- (e) at least one phenol stabilizer (0.05%-0.2% by weight) (e.g., hydroxyquinoline, 0.1%).

Nonionic Detergents. Suitable nonionic detergents include, but are not limited to, at least one selected from the group consisting of adducts of ethylene oxide and fatty alcohols, alkyl phenols, and fatty acid amides, N,N-bis(3-D-gluconamidopropyl)cholamides (BIGCHAP), decanoyl-N-methylglucamides, n-decyl α -D-glucopyranosides, n-decyl β -D-glucopyranosides, n-decyl β -D-maltopyranosides, deoxy-BIGCHAPs, digitonins, n-dodecyl β -D-glucopyranosides, n-dodecyl α -D-maltosides, n-dodecyl β -D-maltosides, heptanoyl-N-methylglucamides, n-heptyl β -D-glucopyranosides, N-heptyl β -D-thiogluconopyranosides, n-hexyl β -D-glucopyranosides, 1-monooleoyl-rac-glycerols, nonanoyl-N-methylglucamides, n-nonyl α -D-glucopyranosides, n-nonyl β -D-glucopyranosides, octanoyl-N-methylglucamides, n-octyl α -D-glucopyranosides, n-octyl β -D-glucopyranosides, octyl β -D-thiogalactopyranosides, octyl β -D-thiogluconopyranosides, polyoxyethylene esters, polyoxyethylene ethers, octylphenoxypoly(oxyethylene)ethanol (IGEPAL CA-360, Triton X-100), polyoxyethylenesorbitan esters (Tween 20), sorbitan esters, tergitols, n-tetradecyl β -D-maltosides, tritons, tyloxapol, and n-undecyl β -D-glucopyranosides. Tritons can include, but are not limited to, triton X-100 (t-octylphenoxypolyethoxyethanol); triton X-100, peroxide- and carbonyl-free; triton X-100, reduced; triton X-100, reduced, peroxide- and carbonyl-free; triton X-114, triton X-405, triton N-101, triton X-405, reduced; other tritons, such as but not limited to the following:


Non-ionic	Low-foam
N-42	CF-10
N-57	CF-21
N-60	CF-32 (95%)
X-15	CF-54
X-35	DF-12
X-45	DF-16
X-102	

Non-ionic	Low-foam
X-155	
X-165 (70%)	
X-207	
X-305 (70%)	
X-705-70 (70%)	
B-1956 (77%)	

Additional suitable nonionic detergents include triton CG-110, triton XL-80N, triton WR-1339 or tyloxapol, tert-octylphenoxy poly(oxyethylene) ethanol (*e.g.*, IGEPAL (Rhône-Poulenc, Paris, France)), and Nonidet P40 (octylphenoxy polyethoxy ethanol).

Chelators. Chelators include, but are not limited to: EDTAs, EGTA's, sodium citrates (citric acids), salicylic acids (and their salts), phthalic acids, 2,4-pentanediones, histidines, histidinol dihydrochlorides, 8-hydroxyquinolines, 8-hydroxyquinoline citrates, and *o*-hydroxyquinones.

Phenols. Any suitable phenolic compound can be used, *e.g.*, according to the following formula I:



where R_1 , R_2 , R_3 , R_4 , R_5 are each independently selected from H, alkyl, halo, *o*-alkyl, acyl and hydroxyl. Examples include, but are not limited to: phenol, *o*-cresol, *m*-cresol, *p*-cresol, resorcinol, β -resorcyraldehyde and the like.

Phenol Stabilizers. Phenol stabilizers include, but are not limited to: 8-hydroxyquinolines, 8-hydroxyquinoline citrates, 2,5,7,8-tetramethyl-2-(4',8',12'-

trimethyltridecyl)-6-chromanols, *p*-hydroxyquinones, *o*-hydroxyquinones, citric acids (and their salts), salicylic acids, ascorbic acids, *p*-phenylenediamines, *n*-propylgallates, and other known radical scavengers.

Phenol Solubilizers. Phenol solubilizers include, but not limited to: any alcohol miscible with phenol and water, *e.g.*, monoalcohols (such as, but not limited to: methyl alcohol, ethyl alcohol, propyl alcohol, and the like); diols (such as, but not limited to: ethylene glycol, propanediol, and the like); and, polyols (such as, but not limited to: glycerol, polyethylene glycol, polyvinyl alcohol, and the like).

The above compounds are commercially available, *e.g.*, from Sigma-Aldrich (St. Louis, MO).

See, e.g., Myers, *Surfactant Science and Technology*, VCH Publishers, Inc., New York (1992); Cutler, *supra*; Rosen, *Surfactants and Interfacial Phenomena*, Wiley, New York (1984); Schick *et al.* *Surfactant Science Series*, Vols. 1-22, Dekker, New York (1961-1987); Tadros, *Surfactants*, Academic Press, London (1984); which references are entirely incorporated herein by reference with regard to their teaching of detergents and/or surfactants.

RNA Recovery Process

An example of a process according to the present invention for providing RNA, includes, but is not limited to:

- (a) mixing the sample with the extraction reagent to form a mixture;
- (b) adding a haloalkane to the mixture;
- (c) separating the organic and aqueous phases; and
- (d) precipitating the cytoplasmic RNA from the aqueous phase obtained in step (c).

The process can optionally further comprise: (e) recovering the cytoplasmic RNA from the precipitate obtained in step (d). The process can also optionally further comprise: (f) isolating mRNA from the recovered cytoplasmic RNA precipitate

using any known method, such as by chromatography on oligo(dT)-cellulose (see, e.g., Sambrook, *infra*, at § 7.26-7.29).

5 In step (b) of the above method, the haloalkane added to the sample and extraction reagent mixture can be any haloalkane suitable for separating RNA from other cytoplasmic components. Such haloalkanes include, but are not limited to chloroform, methylene chloride (dichloromethane), 1-bromo-3-chloropropane, 2-bromo-1-chloropropane, bromoethane, 1-bromo-5-chloropentane, and bromotoluene.

10 In step (c) of the above method, the phases can be separated using any suitable method for separating RNA in an aqueous phase from other cytoplasmic components in the organic phase. Such separation methods include, but are not limited to centrifugation, filtration, vacuum filtration, or gravity.

15 In step (d) of the above method, the cytoplasmic RNA can be separated using any suitable method for precipitating RNA from the aqueous phase. Such separation methods include, but are not limited to using alcohol, polyethylene glycol, lithium chloride, or the like.

20 In step (e) of the above method, the cytoplasmic RNA can be recovered using any suitable method for recovering RNA from the precipitate. Such separation methods include, but are not limited to dissolving the RNA in water or in low salt buffer.

Having now generally described the invention, the same will be more readily understood through reference to the following example which is provided by way of illustration, and is not intended to be limiting of the present invention.

Example 1: Isolation of Plant RNA

Protocol for Isolation of Cytoplasmic RNA from a Plant Sample or Specimen

25 The plant specimen is ground into a powder in liquid nitrogen. The plant powder is stored and used frozen. 0.1 g of the frozen powdered plant is thoroughly mixed with 1 ml of the RNA extraction reagent of the invention and

is then let stand at room temperature for 5 minutes. 0.2 ml of chloroform per ml of the RNA extraction reagent is added to the mixture, and mixed thoroughly. The sample is centrifuged for 5 minutes at 12,000 x g at room temperature. (Larger quantities can be centrifuged for 30 minutes at 2600 x g or 6000 x g for 10 minutes at 4°C.)

The aqueous phase is transferred to an RNase-free tube. 0.5 ml of isopropanol per ml of RNA extraction reagent is added. The sample is mixed and let stand at room temperature for 10 minutes. The sample is centrifuged for 10 minutes at 12,000 x g at 4°C. (Larger quantities can be centrifuged for 30 minutes at 26000 x g or 6000 x g for 10 minutes at 4°C.) The supernate is decanted. The pellet is then washed with 1 ml of 75% ethanol per ml of RNA extraction reagent, centrifuged and any residual liquid is removed. RNase-free water is added to dissolve the RNA (50 µl of water for the 0.1 g sample).

Isolation of cytoplasmic RNA from cells grown in suspension. The cells are collected by centrifugation at 4°C for 5 minutes at 2000 g. The fresh or frozen cell pellet containing 1×10^7 cells is gently resuspended in 1 ml of RNA extraction reagent. The mixture is allowed to stand at room temperature for 5 minutes.

0.2 ml of chloroform is added per ml of RNA extraction reagent used., and mixed thoroughly. The sample is centrifuged for 5 minutes at 12,000 x g at room temperature. (Larger quantities can be centrifuged for 30 minutes at 2600 x g or 6000 x g for 10 minutes at 4°C.) The top, aqueous phase is transferred to an RNase-free tube, and 0.5 ml of isopropanol is added per ml of RNA extraction reagent. The supernatant is mixed and let stand at room temperature for 10 minutes.

The sample is centrifuged for 10 minutes at 12,000 x g at 4°C. (Larger quantities can be centrifuged for 30 minutes at 2600 x g or 6000 x g for 10 minutes at 4°C.) The supernate is decanted. It is then washed with 1 ml of 75% ethanol per ml of RNA extraction reagent, centrifuged and any residual liquid is removed. RNase-free water is added to dissolve the RNA (50 µl of water for the 1×10^7 cells sample).

The concentration of the RNA is determined by measuring the OD_{260} of an aliquot of the final preparation. A solution of RNA whose $OD_{260} = 1$ contains approximately 40 μ g of RNA per milliliter.

Results using the above Protocol.

5 The following Table 1 shows results obtained using the above protocol, as RNA yields.

Table 1

Sample	Method	Amount (g)	Ratio $A_{260/280}$	RNA (μ g)
Tobacco leaves (soil grown plant)	RNA reagent	0.10	2.0	15
Tobacco leaves (soil grown plant)	RNA reagent	1.0	1.7	212
Tobacco leaves (soil grown plant)	RNA reagent	1.0	1.8	191
Tobacco leaves (soil grown plant)	Trizol	1.0	1.8	181
Tomato plant (culture)	RNA reagent	0.11	1.9	38
Tomato plant (culture)	RNA reagent	1.0	1.9	351
Tomato plant (culture)	Trizol	0.8	1.7	295
Blue spruce needles (dormant tree)	RNA reagent	0.11	1.5	16
Blue spruce needles (dormant tree)	RNA reagent	2.7	1.4	264
Blue spruce needles (dormant tree)	Trizol	2.7	0.6	12
Grass	RNA reagent	1.0	1.8	463
Grass	Trizol	1.0	1.8	160
Fungal hyphae, sense	RNA reagent	10^9 cells	1.8	274
Fungal hyphae, anti-sense	RNA reagent	10^9 cells	1.9	57
HeLa cells	RNA reagent	10^8 cells	1.8	2296

Having now fully described this invention it will be understood to those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are entirely incorporated by reference herein.

5

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

What Is Claimed Is:

1. An aqueous RNA isolation reagent, comprising two or more components selected from the group consisting of

- (a) at least one nonionic detergent at a concentration of 0.1-1.0% (vol/vol);
- (b) at least one chelator at a concentration of 0.02-0.25 M; and
- (c) at least one phenol at a concentration of 10%-60% (wgt/vol); and
- (d) at least one phenol solubilizer at a concentration of 15%-55% (vol/vol).

2. An RNA isolation reagent according to claim 1, wherein said at least one nonionic detergent is present at a concentration of 0.5-0.8% (vol/vol).


3. An RNA isolation reagent according to claim 3, wherein said nonionic detergent comprises at least one detergent selected from the group consisting of an octylphenoxypoly(oxyethylene)ethanol, an N,N-bis(3-D-glucanamidopropyl)cholamide (BIGCHAP), an decanoyl-N-methylglucamide, an n-decyl α -D-glucopyranoside, an n-decyl β -D-glucopyranoside, an n-decyl β -D-maltopyranoside, a deoxy-BIGCHAP, a digitonin, an n-dodecyl β -D-glucopyranoside, an n-dodecyl α -D-maltoside, an n-dodecyl β -D-maltoside, a heptanoyl-N-methylglucamide, an n-heptyl β -D-glucopyranoside, an N-heptyl β -D-thioglucopyranoside, an n-hexyl β -D-glucopyranoside, a 1-monooleoyl-rac-glycerol, a nonanoyl-N-methylglucamide, an n-nonyl α -D-glucopyranoside, an n-nonyl β -D-glucopyranoside, an octanoyl-N-methylglucamide, an n-octyl α -D-glucopyranoside, an n-octyl β -D-glucopyranoside, an octyl β -D-thiogalactopyranoside, an octyl β -D-thioglucopyranoside, a polyoxyethylene ester, a polyoxyethylene ether, a polyoxyethylenesorbitan ester, Tween 20, a sorbitan ester, an n-tetradecyl β -D-maltoside, a triton, a tyloxaapol and an n-undecyl β -D-glucopyranoside.

4. An RNA extraction reagent according to claim 3, wherein said non-ionic detergent is a octylphenoxypoly(oxyethylene)ethanol.

5. An RNA extraction reagent according to claim 4, wherein said octylphenoxy(poly(oxyethylene))ethanol is present at a concentration of about 0.5% (vol/vol).

6. An RNA extraction reagent according to claim 1, wherein said chelator is selected from the group consisting of sodium citrate, EDTA, EGTA, sodium citrate, a citric acid, a salicylic acid or a salt thereof, a tergitol, a phthalic acid, a 2,4 pentanedione, a histidine, a histidinol dihydrochloride, an 8-hydroxyquinoline, an 8-hydroxyquinoline citrate and an *o*-hydroxyquinone.

7. An RNA extraction reagent according to claim 1, wherein said phenol is a compound according to formula I:



where R_1 , R_2 , R_3 , R_4 , R_5 are each independently selected from H, alkyl, *o*-alkyl, halo, acyl and hydroxyl.

8. An RNA extraction reagent according to claim 1, further comprising at least one phenol solubilizer at a concentration of 15%-55% (wgt/vol).

9. An RNA extraction reagent according to claim 8, wherein said phenol solubilizer is selected from the group consisting of a monoalcohol, a diol and a polyol.

5 10. An RNA extraction reagent according to claim 1, further comprising at least one phenol stabilizer at a concentration of 0.05%-0.2% (wgt/vol).

10 11. An RNA extraction reagent according to claim 10, wherein said phenol stabilizer is at least one selected from the group consisting of hydroxyquinoline, 8-hydroxyquinoline, 8-hydroxyquinoline citrate, 2,5,7,8-tetramethyl-2-(4',8',12'-trimethyltridecyl)-6-chroman-1-ol, *p*-hydroxyquinone, *o*-hydroxyquinone, citric acid or salt thereof, salicylic acid, ascorbic acid, *p*-phenylenediamine and *n*-propylgallate.

15 12. A kit for isolation of RNA, comprising at least one container, wherein a first container contains at least one RNA extraction reagent according to claim 1.

13. A method for providing cytoplasmic RNA from a sample comprising eukaryotic cells, said method comprising

- 20 (a) mixing said sample containing said cells with an RNA extraction reagent according to claim 1 to form a mixture;
- (b) adding a haloalkane to the mixture and mixing the resulting organic and aqueous phases;
- (c) separating the organic and aqueous phases; and
- (d) precipitating cytoplasmic RNA from the aqueous phase obtained in step (c).

14. A method according to claim 13, further comprising
(e) recovering the cytoplasmic RNA from the precipitate obtained in step (d).

5

15. A method according to claim 14, further comprising
(f) isolating mRNA from said cytoplasmic RNA.

16. A method according to claim 14, wherein said sample is derived from a plant or a plant material.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
220

RNA Isolation Reagent and Methods

Abstract


The invention provides methods, extraction reagents and kits for RNA isolation from eukaryotic cells such as plant or animal cells, where a phenol, a phenol solubilizer, a chelator and a nonionic detergent are used in the extraction reagent to replace the need for chaotropes and/or RNase inhibitors.

5

a40-04.wpd

United States Patent & Trademark Office

Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

1. Application papers are not suitable for scanning and are not in compliance with 37 CFR 1.52 because:
 - ☐ All sheets must be the same size and either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
Pages _____ do not meet these requirements.
 - ☐ Papers are not flexible, strong, smooth, non-shiny, durable, and white.
 - ☐ Papers are not typewritten or mechanically printed in permanent ink on one side.
 - ☐ Papers contain improper margins. Each sheet must have a left margin of at least 2.5 cm (1") and top, bottom and right margins of at least 2.0 cm (3/4").
 - ☐ Papers contain hand lettering.
2. Drawings are not in compliance and were not scanned because:
 - ☐ The drawings or copy of drawings are not suitable for electronic reproduction.
 - ☐ All drawings sheets are not the same size. Pages must be either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
 - ☐ Each sheet must include a top and left margin of at least 2.5 cm (1"), a right margin of at least 1.5 cm (9/16") and a bottom margin of at least 1.0 cm (3/8").
3. Page(s) _____ are not of sufficient clarity, contrast and quality for electronic reproduction.
4. Page(s) _____ are missing.
5. OTHER: NO DECLARATION OF Drawing